

GENERAL INDEX TO VOLUME XXXVII

New scientific names of plants and the final members of new combinations are printed in **bold-face** type; synonyms and page numbers having reference to figures and plates, in *italics*; and all other matter in ordinary type.

A

- Acitheca, 418, 424, 446, 455, 457
- Acnida, 610
- Adipera santanderensis, 402
- Afghanistan, cultivation and use of grain amaranths in, 593
- Africa, cultivation and use of grain amaranths in, 594
- Alafia Schumannii*, 406
- Albizia coripatense, 400; paucipinnata, 400
- Alegria manufacture from grain amaranths, 578
- Amaranths, The Grain: A survey of their history and classification, 561; Appendix (Extracts from early Latin American Sources), 629
- Amaranths, grain: alegria manufacture from, 578; botanical classification, 594; chromosome counts in, 596; cultivated species, 597; morphology of, 596, 598, 604-605; cultivation and use, 562, in Africa, 594, in Asia, 588, in Guatemala, 582, 583, in Mexico, 564, 574, 575, 578, 625, in the United States, 563; huauhtli, in relation to, 567, 624; niguas, made from, 584, 626; non-cultivated species, 603; pistillate flowers of, 598, 604-605
- Amaranthus: *Alopecurus*, 594, 602; sect. *Amaranthophytus*, 594, 612; *Anardana*, 590, 594, 599; *ataco*, 612; *Bigelovii*, 606; *bracteosus*, 606; *Brandegei*, 606; *caudatus*, 585, 588, 594, 602, 628; *celosioides*, 594, 608; *chihuahuensis*, 612; *cruentus*, 594, 601, common race, 601, 627; Mexican race, 601, 628; cultivated species, 597; *dubius*, 607; *edulis*, 588, 594, 597, 603, 628; *farinaceus*, 594; *flavus*, 594; *frumentaceus*, 589, 594, 599; *hybridus*, 594, 608, northern race, 609, tropical race, 610; *hypochondriaceus*, 594, 599, 609; *leucocarpus*, 563, 594, 599, 610, aberrant race, 600, 627, Arizona race, 600, 627, common race, 599, 627; *leucospermus*, 594, 599; non-cultivated species, 603; *paniculatus*, 594, 599; *Powellii*, 594, 611, Cordilleran race, 612, Great Basin race, 611; *quitensis*, 608; *retroflexus*, 594, 605; *sanguineus*, 594; *scariosus*, 607; *speciosus*, 594; *spinosis*, 605; *strictus*, 594; *viscidulus*, 606; *Wrightii*, 612
- Amblystigma pedunculare*, 540
- Ampbistelma exsertum*, 521
- Anachoropteris, 450, 452
- Angiopteris, 409, 447, sori of, 454
- Ankyropteris, 412
- Aphonostelma micranthum, 552
- Apocynaceae, 404, 496
- Archaeological finds of grain amaranth seeds, 563, 587
- Archangiopteris, 444, 454
- Argentina, cultivation and use of grain amaranths in, 587
- Arizona, grain amaranths from, 563
- Asclepiadaceae, 496, 503; (*Miscellaneae*), 406; adaptation of, to insect pollination, 477; taxonomy of, 503
- Asclepias angustissima*, 535; *viminalis*, 511
- Asia, cultivation of grain amaranths in, 588; regions where indicated, 591
- Asterotheca, 418, 455, 457
- Asterohceceae, 418, 446; showing two possible lines of derivation of, 455
- Ateleia, 401
- Atriplex, 564
- Aztecs, use of grain amaranths by, 564, 569

B

- Bailey, L. H. *Rubus* in Panama, 154
- Bartramia*, 317, 332; *indica*, 317, 382
- Bledo, 584
- Blepharodon, 504; *mucronatum*, 504, 552; *nitidus*, 504
- Blitum, 564
- Blitum majus Peruanum, 585
- Bolivia, cultivation and use of grain amaranths in, 585, 586
- Botrychoxylon, 412
- Botryopteris, 413, 452; *americana*, 413; *antiqua*, 413, 416; *dubius*, 413; *fecunda*, 414, 415, 460; *forensis*, 410, 413, 414; *globosa*, 410, 414; *hirsuta*, 413, 416; *illinoensis*, 415, 462; *ramosa*, 413, 416; *spinosa*, 415, 460, 462

C

- Carboniferous fern fructifications, Some American, 409

- Cerica dolichaula*, 404
Caricaceae, 404
Cassia tomentosa, 402; *vargasi*, 401
Ceramantbus, 480, 505, 506; *flavus*, 536; *gracilis*, 521
 Ceylon, grain amaranths in, 589
Chansitheca, 443
Chenopodium, 564; *Nuttallii*, 565; *puebloense*, 565; *quinoa*, 566
 China, grain amaranths in, 589, 593
Chlaenaceae, pollen of, 497
Chorionopteris, 412, 446, 450, 455, 457; *gleichenioides*, 449, 450, 451, 452; *sporangium* of, 448, 449, 450
Christensenia, 447, 457
 Coal Age plants, 409
Coenopteridales, 412, soral morphology of, 448, 449
Connaraceae of Panama, 178
Corona of *Sarcostemma*, 491
Corynepteris, 446, 448; *Essenghi*, 449; *coralloides*, 449
Crassulaceae of Panama, 121
Cuahzontli, 566
Cunoniaceae of Panama, 145
Cyathostegia, 401; *matthewsii*, 401; *weberbaueri*, 401
Cyathotrichus, 419, 455; *altissimus*, 420, 464; *altus*, 419; *bulbaceus*, 419
Cynanchum, 504; *clausum*, 511; *mexicanum*, 511
Cystostemma, 479, 505, 506; *umbellatum*, 480, 536
- D
- Danaea*, 443, 454
Danaeites, 443
Danaeopsis, 444
Doemia, 481
- E
- Economic use of huauhtli, 572
 England, coal balls from, 412
Eoangiopterus Andrewsii, 440, 476; reconstruction of, 442; sori of, 441, 453, 454
Epteris, 448, 449
 Ethnological problems, grain amaranths in relation to, 561
 Evolutionary possibilities within Marattiaceae, 452
- F
- Fern fructifications, Some American carboniferous, 409
Fischeria, 492
 Flora of Panama. Part IV, Fasc. 1, 1; Part V, Fasc. 2, 121
 Fossil plants, 410
 Fructifications, fern, 409
- Funastrum*, 479, 505, 506; *angustissimum*, 479, 535; *apiculatum*, 511; *barbatum*, 511; *bicolor*, 525; *bilobum*, 517; *bonariense*, 511; *clausum*, 512; *crassifolium*, 512; *crispum*, 522; *cumanense*, 512; *cupidatum*, 512; *cynanchoides*, 528; *Dombeyanum*, 512; *dumetorum*, 519; *elegans*, 525; *Eruvendbergii*, 519; *Fendleri*, 519; *flavum*, 536; *fragile*, 512; *Gardneri*, 512; *glaucum*, 509; *Glaziovii*, 512; *gracile*, 521; *Hartwegii*, 530; *heterophyllum*, 523; *birtellum*, 532; *lanceolatum*, 512; *hsianthum*, 512; *Lindenianum*, 519; *lineare*, 530, var. *heterophyllum*, 530; *longifolium*, 536; *luridum*, 525; *odoratum*, 520; *ovalifolium*, 509; *pallidum*, 512; *Palmeri*, 512; *pannosum*, 507; *Pavoni*, 507; *pedunculatum*, 512; *pubescens*, 512; *reflexum*, 519; *refractum*, 517; *riparium*, 512; *rotundifolium*, 507; *Schottii*, 512; *Sebertii*, 512; *tomentellum*, 507; *Torreysi*, 527; *vesiculare*, 527
- G
- Galactia*, 402; *gracillima*, 403; *linearifolia*, 403; *pedunculata*, 402
- Gonolobus*, 492
- Grain amaranths, 561
- Gramineae* (*Miscellaneae*), 397
- Grand'eurya*, 443
- Guatemala, cultivation of grain amaranths in, 582
- H
- Helicocarpus*, 316, 326; *arborescens*, 344; *birtus*, 374; *occidentalis*, 349; *tigrinus*, 384
- Herreria *Glaziovii*, 398; *latifolia*, 397
- Holalafia multiflora*, 406; *Schumannii*, 406
- Holm, R. W.: The genus *Sarcostemma*, 477
- Huauhtli: botanical identity of, 564; cultivation, 572; economic use, 572; in religious ceremonies, 564; recent cultivation of, 577
- Huauhquilitl, 624
- Huauhzontli, 566
- I
- Illinois, coal balls from, 411
- Inca grain, 585
- India, grain amaranths in, 589
- Indiana, coal balls from, 411
- Indians of the Americas, role of grain amaranths among, 563, 564, 585
- Iowa, coal balls from, 411
- Inga alatopetiolata*, 206; *edulis* var. *minutula*, 209; *gigantifolia*, 218; *panamensis* var.

INDEX

- clavata, 204, var. *Pittieri*, 203, var. *Rodrigueziana*, 205
- Insect pollination: in Asclepiadaceae, 477; in Orchidaceae, 477, 497
- J
- Jacaratia dolichaula*, 404
- K
- Kansas, coal balls from, 411
- Keramanthus*, 480
- L
- Lay, Ko Ko. The American species of *Triumfetta* L., 315
- Leguminosae: (*Miscellaneae*), 398; of Panama, 184; Subfamily Mimosoideae, 185, 497
- Leucaena multicapitula*, 302
- Liliaceae (*Miscellaneae*), 397
- Lochnera rosea*, 491
- Loganiaceae (*Miscellaneae*), 404
- Lugonia*, 481, 505, 537; *andina*, 549; *lysimachioidea*, 547
- M
- McClintock, Elizabeth. *Saxifragaceae* of Panama, 137
- McVaugh, Rogers. *Rosaceae* of Panama, 147
- Macroglossum*, 444, 454
- Mamay, Sergius Harry. Some American carboniferous fern fructifications, 409
- Manchuria, grain amaranths in, 593
- Mandevilla *Espinosa*, 405; *glandulosa*, 406; *jasminiflora*, 405; *ligustriflora*, 404; *scutifolia*, 406; *tricolor*, 406
- Marathrum cheiriferum*, 134; *indifferens*, 132; *leptophyllum*, 131; *pussillum*, 129; *stenocephalum*, 127
- Marattiaceae, 409; evolutionary possibilities among, 452, 453; fossil, 445, fructifications compared with pteridospermous microsporangiate organs, 446; living, 444, 447; origin of, 444, 447; phyletic chart of, 450; phyletic slide of, 445, 456; sori of, 418, 448, 453, linear, 452, radial, 457
- Marattiales, 409, 418
- Marattiopsis, 444
- Marsdenia *Dussii*, 408; *Rotheana*, 408
- Matelea *belizensis*, 407; *cyclophylla*, 408; *Gentlei*, 407; *ionantha*, 408; *Meyeri*, 406; *reticulata*, 407; *Rogersii*, 407
- Mexhuauhtli, 565
- Mexico, cultivation and use of grain amaranths in, 564; localities where indicated, 574, 575
- Michihuauhtli, 562, 624
- Mimosoideae (subfamily of Leguminosae), 185, 497
- Miscellanea taxonomica I*, 397
- Moringaceae* of Panama, 121
- N
- Nexhuauhtli, 565
- Niguas, manufacture of, from grain amaranths, 584; popped amaranth seed prepared for, 626; made in Guatemala, 626
- O
- Oligocarpia*, 443
- Ophioglossaceae*, 445
- Orchidaceae, adaptation of, to insect pollination, 477, 497
- Ottonia Thiemeana*, 71
- Oxystelma*, 480, 503, 537; *Gilliesii*, 540; *solanoides*, 545; *Vaileae*, 543; *violacea*, 548
- P
- Paleobotany, aim of, 444
- Panama: Flora of, Part IV, Fascicle 1 (*Piperaceae*), 1; Part V, Fascicle 2 (Resedaceae to Leguminosae, in part), 121
- Paracopteris*, 443
- Pari*, 580
- Pecopteris*, 409, 447; *intermedia*, 418; polymorpha, 424
- Pentacyphus*, 505, 538, 548; *boliviensis*, 549
- Pentagonium*, 481, 505, 537; *flavum*, 544
- Peperomia caudilimba* var. *cylindribacca*, 104; *chiriquiensis*, 93; *Chrysleri*, 94; *conserta*, 114; *Davidsonii*, 102; *flexinervia*, 113; *glabella* var. *nervulosa*, 98; *hispidorrhachis*, 112; *oblongibacca* var. *macrophylla*, 105, var. *marragantina*, 105, var. *subvillosa*, 106; *obtusifolia* var. *longibracteata*, 109; *perglandulosa*, 116; *umbigaudens*, 81; *Wedelii*, 118
- Pergularia*, 481
- Periploca*, 492; *esculenta*, 482
- Peru, cultivation and use of grain amaranths in, 585; localities where indicated, 586
- Petziat*, 568, 624
- Philibertia*, 480, 505, 506; *crispa*, 522; *clausa*, 511; *crassifolia*, 511; *cumanensis*, 511; *cynanchoides*, 528; *elegans*, 525; *filipes*, 519; *Hartwegii*, 522, 530, var. *beterophylla*, 522, 530; *beterophylla*, 522, 530; *birtella*, 532; *lasiantha*, 511; *ovalifolia*, 509; *pallida*, 511; *Palmeri*, 511; *pedunculata*, 511; *riparia*, 511; *Torreyi*, 527
- Philibertia*, 479, 505, 506, 537; *albiflora*, 543; *anomala*, 552; *bicolor*, 525; *biloba*, 517; *bonariensis*, 511; *campanulata*, 539; *clausa*, 511; *crassifolia*, 511; *crispa*, 522; *cumanensis*, 511; *cuspidata*, 511; *cynanchoides*, 528, var. *subtruncatum*, 528; *dume-*

- torum*, 519; *elegans*, 525; *Ervendbergii*, 519; *exserta*, 521; *Fendleri*, 519; *flava*, 544; *Gardneri*, 511; *Gilliesii*, 540, var. *albida*, 541, var. *gracilis*, 541, var. *grandiflora*, 541, var. *mirabilis*, 541, var. *parviflora*, 541, var. *pubescens*, 541, var. *vestita*, 541; *gracilis*, 540; *grandiflora*, 540, f. *albida*, 541; *beterophylla*, 523, 530; *birtella*, 532; *hypoleuca*, 539; *Lindeniana*, 519; *linearis*, 530, var. *heterophylla*, 522, 530, var. *birtella*, 532; *longifolia*, 536; *lurida*, 525; *lysimachiooides*, 547; *obtusata*, 545; *obtusiuscula*, 545; *odorata*, 520; *Palmeri*, 511; *pannosa*, 507; *Pavoni*, 507; *picta*, 544; *quadriflora*, 545; *reflexa*, 519; *refracta*, 517; *riparia*, 511; *rotundifolia*, 507; *scandens*, 552; *Schreiteri*, 541; *solanooides*, 544; *speciosa*, 539; *splendens*, 539; *stellaris*, 536; *stipitata*, 546; *tomentella*, 507; *Torreyi*, 527; *undulata*, 522; *varifolia*, 545; *viminalis*, 511; *violacea*, 541; *viridiflora*, 552; *Web-erbaueri*, 545
- "Phyletic slide" of Marattiaceae, 445, 450, 451
- Phylogeny of ferns, 444, 456
- Piper acutissimum* var. *trichopus*, 66; *aduncum* var. *brachyarthrum*, 32; *aequale* var. *laurifolium*, 48; *arboreum* var. *falcifolium*, 65, var. *hirtellum*, 64; *aristochiaeifolium*, 18; *augustum* var. *coelestanum*, 61; *campanum*, 25; *Cooperi*, 21; *crassinervium* var. *tocotanum*, 8; *Daviddsonii*, 14; *dimorphotrichum*, 52; *distigmatum*, 13; *hastularum*, 21; *hirtellipetiolatum* var. *Harveyanum*, 58, var. *taianum*, 58; *hispidum* var. *ellipticifolium*, 34, var. *trachydermum*, 33; *lucigaudens* var. *Allenii*, 42; *macropunctatum*, 10; *marginatum* var. *clausum*, 17; *Maxonii* var. *varium*, 71; *perbrevicaule*, 51, var. *subglabrum*, 51; *perlaseum*, 67; *septoplirnervium* var. *subpubinervium*, 46; *subcaudatum* var. *maternale*, 59; *variitrichum*, 37; *via-chicoense*, 67; *Wedelii*, 56, var. *retrohirsutum*, 57
- Piperaceae of Panama, 1
- Pithecellobium*, 398; *coripatense*, 400; *membraraceum*, 228; *multiflorum*, 400; *ramiflorum*, 400; *steyermarkii*, 398
- Podostemoneaceae of Panama, 124
- Pollination, insect: in Asclepiadaceae, 477; in Orchidaceae, 477, 497
- Polypodiaceae, 445
- Psaronius*, 409, 447, 452
- Pteridospermae, 409
- Pteridospermous microsporangiate organs and marattiaceous fructifications compared, 446
- Ptychoscarpus*, 418, 423, 447, 455; *unitus*, 418, 454, 457, 462
- Q
- Quihuica*, 585
- Quelite*, 580
- Quinna*, 585
- Quinoa*, 585
- R
- Ramirez Codex*, 567
- Relaciones of Mexico, 576
- Religious festivals, use of grain amaranths in, 564
- Resedaceae of Panama, 121
- Rhynia*, 448
- Rinorea roureoides*, 403
- Rosaceae of Panama, 147
- Rourea glabra*, 404
- Rubus in Panama, 154
- S
- Sarcostemma, R. Br., (Asclepiadaceae), The American species of, 477
- Sarcostemma, 479, 482, 537, 538; *andinum*, 549; *angustissimum*, 534, 535; *apiculatum*, 511; *arenarium*, 533; *barbatum*, 511; *bellum*, 517; *bicolor*, 525; *bifidum*, 511; *bilobum*, 517, ssp. *bilobum*, 517, ssp. *Lindenianum*, 517, 519; *bilobum*, 517; *bonariense*, 511; *Brownei*, 511; *campanulatum*, 538, 539; *carpophyloides*, 521; *clausum*, 509, 511; *crassifolium*, 511; *crispum*, 522; *cumanense*, 511; *cuspidatum*, 511; *cyanchooides*, 516, 528, ssp. *cyanchooides*, 528, ssp. *Hartwegii*, 528, 530, cl. *cyanchooides-Hartwegi*, 532; *cyanchooides*, 525; *Dombeyanum*, 511; *Donianum*, 520; *elegans*, 506, 525; *esculentum*, 482, 537; *flavum*, 536; *Gardneri*, 511; *Gilliesii*, 540; *glaucum*, 509; *Glaziovii*, 511; *gracile*, 521; *grandiflorum*, 540; *bastatum*, 545; *beterophyllum*, 522, 530, var. *hirtellum*, 532; *hirtellum*, 532; *incanum*, 540; *Jacquinii*, 551; *Lehmanni*, 550; *Lindenianum*, 519; *lineare*, 511, 530; *lobata*, 523; *luridum*, 525; *lysimachiooides*, 538, 547; *marsupiflorum*, 545; *odoratum*, 520; *pallidum*, 511; *pannosum*, 507; *pedunculatum*, 511; *pubescens*, 511; *quadriflorum*, 545; *riparium*, 511; *rotundifolium*, 501; *Schottii*, 511; *solanooides*, 538, 544; *stipitatum*, 546; *Swartzianum*, 511; *tomentellum*, 507; *Torreyi*, 527; *trichopetalum*, 551; *undulatum*, 522; *Vaileae*, 543; *variifolium*, 545; *viminale*, 506
- Sauer, Jonathan Deininger. The grain amaranths: A survey of their history and

- classification, 561
Saxifragaceae of Panama, 137
 Schery, Robert W. *Miscellanea Taxonomica*: Leguminosae, 398; Leguminosae of Panama, 184; Robert E. Woodson, Jr., and collaborators. *Flora of Panama*, Part IV, Fasc. 1, 1; Part V, Fasc. 2, 121
Schizaeaceae, 451
Scolecopteris, 410, 423, 446, 450, 451, 455; comparison of various species of, 438; *elegans*, 425; *incisifolia*, 433, 435, 450, 472; *iowensis*, 429, 454, 468; *latifolia*, 425, 466; *major*, 436, 474; *minor*, 424, var. *parvifolia*, 431, 470; *Oliveri*, 425, 436; *polymorpha*, 424; *Radforthii*, 425; *ripariensis*, 424; *subelegans*, 424
Sori of *Marattiales*, 418, 448, 449; derivation of linear, 452, of radial, 457
 South America, grain amaranths in, 585; localities of, 586
Spigelia aceifolia, 404; *stenophylla*, 404
Sporobolus hians, 397; *pseudairoides*, 397
Stapelia, 492
Stauropteris, 412, 446, 450; *Oldhamia*, 449
Stipitopteris, 452
Sturiella, 418
Swartzia Sect. *Cyatostegia*, 401
- T
- Tetraphysa*, 481, 505, 537; *Lehmannii*, 550
Tezchahuauhtli, 565, 568, 624
Tlapalhuauhtli, 565, 624
Triumfetta, The American species of, 315; chromosome counts, 332; generic and specific relationships, 326; geography, 318; history, 317; morphology, 322; taxonomy, 322
Triumfetta, 332; *abutiloides*, 365; *acrantha*, 370; *acuminata*, 357; *acutiloba*, 373; *althaeoides*, 371; *angulata*, 382; *apetala*, 385; *arboreascens*, 344; *attenuata*, 347; *barbosa*, 360; *Bartramia*, 382; *Berteri*, 378; *bogotensis*, 366; *Botteriana*, 367; *boyacana*, 363; *brachistacantha*, 381; *brachypetalia*, 388; *brevipes*, 352; *Calderoni*, 343; *calycina*, 388; *caudata*, 341; *chihuahuensis*, 358; *columnaris*, 362; *columnarioides*, 362; *coriacea*, 340; *cucullata*, 339; *cymosa*, 363, var. *glabrescens*, 363, var. *birsuta*, 363; *debiscens*, 364; *dichotoma*, 371; *dioica*, 347; *discolor*, 364; *dumetorum*, 367; *eriocarpa*, 382; *excisa*, 382; *falcifera*, 355; *ferruginea*, 344; *Galeottiana*, 385; *Goldmani*, 340; *grandiflora*, 363, var. *brasiliensis*, 363; *grossulariaefolia*, 367; *bavanensis*, 373; *heliocarpoides*, 388; *heterophylla*, 378; *Hintonii*, 342; *birta*, 373; *bispida*, 367; *Hostmanni*, 378; *insignis*, 337; *josefina*, 367; *Jelskii*, 388; *Lappula*, 378; *Lappula*, 382; *leiocarpa*, 384; *Lindeniana*, 367; *longicoma*, 358; *longicuspis*, 363; *longiseta*, 363; *macrocalyx*, 338; *Martiana*, 382; *Matudai*, 384; *Mexiae*, 348; *micropetala*, 358; *microphylla*, 371; *mollissima*, 357; *multilocularis*, 356; *nemoralis*, 363; *obscura*, 369; *obovata*, 337; *oligacantha*, 377; *Orizaba*, 367; *ovata*, 373; *oxyphylla*, 373; *Palmeri*, 353; *panamensis*, 367; *paniculata*, 384; *persimilis*, 346; *pilos*, 366; *polyandra*, 337; *pseudocolumnaris*, 362; *Purpusii*, 354; *quergetorum*, 340; *quinqueloba*, 378; *rhombeaefolia*, 382; *rhomboidea*, 382, var. *recifensis*, 328, var. *Spruceana*, 382; *rubricaulis*, 373; *Salzmanni*, 378; *Sampaioi*, 384; *sanctae-Luciae*, 340; *Schunkei*, 363; *semiriloba*, 373, f. *althaeoides*, 371, var. *Berlandieri*, 374, var. *brasiliensis*, 374, var. *bavanensis*, 374, var. *Martiana*, 374, 382, var. *surinamensis*, 371, var. *typica*, 374; *sepium*, 367; *sericata*, 380; *sinuosa*, 378; *Sloanei*, 374; *socorensis*, 353; *speciosa*, 338; *stellata*, 334; *surinamensis*, 371; *Thonningiana*, 373; *tigrina*, 384; *tiliaeefolia*, 373; *tricuspidis*, 373; *triloba*, 373; *valenciensis*, 367; *velutina*, 358; *vincentina*, 367
- U
- United States, cultivation and use of grain amaranths in, 563
- V
- Van Royen, P. *Podostemonaceae of Panama*, 124
 Van Schaack, George B. *Gramineae (Miscellaneae)*, 397
Vellozia, 398; *caruncularis*, 398; *graminea*, 398; *Macedonis*, 398
Velloziaceae (Miscellaneae), 398
Violaceae (Miscellaneae), 403
- W
- Weywi, 582
 Woodson, Robert E., Jr. *Miscellanea taxonomica*: *Apocynaceae*, 404; *Asclepiadaceae*, 404; *Caricaceae*, 404; *Liliaceae*, 397; *Loganiaceae*, 404; *Velloziaceae*, 403; *Violaceae*, 403; Robert W. Schery and, and collaborators. *Flora of Panama*, Part IV, Fasc. 1, 1; Part V, Fasc. 2, 121
- X
- Kochihuauhtli, 565, 624
- Y
- Yuncker, T. G. *Piperaceae of Panama*, 1
- Z
- Zoale, 564, 568, 570, 576
Zosima, 481, 537; *violacea*, 540
Zygopterus, 413